



<https://publications.dainst.org>

iDAI.publications

DIGITALE PUBLIKATIONEN DES
DEUTSCHEN ARCHÄOLOGISCHEN INSTITUTS

Das ist eine digitale Ausgabe von / This is a digital edition of

Schwanghart, Wolfgang – Möller, Steffen – Schütt, Brigitta

Environmental Characteristics of the Upper and Middle Orkhon Valley, Mongolia.

in: Bemann, Jan – Érdénébat, Ulambaiaryn – Pohl, Ernst (Hrsg.), Mongolian-German Karakorum expedition. 13–26.

DOI: <https://doi.org/10.34780/133a-8dr5>

Herausgebende Institution / Publisher:
Deutsches Archäologisches Institut

Copyright (Digital Edition) © 2022 Deutsches Archäologisches Institut
Deutsches Archäologisches Institut, Zentrale, Podbielskiallee 69–71, 14195 Berlin, Tel: +49 30 187711-0
Email: info@dainst.de | Web: <https://www.dainst.org>

Nutzungsbedingungen: Mit dem Herunterladen erkennen Sie die Nutzungsbedingungen (<https://publications.dainst.org/terms-of-use>) von iDAI.publications an. Sofern in dem Dokument nichts anderes ausdrücklich vermerkt ist, gelten folgende Nutzungsbedingungen: Die Nutzung der Inhalte ist ausschließlich privaten Nutzerinnen / Nutzern für den eigenen wissenschaftlichen und sonstigen privaten Gebrauch gestattet. Sämtliche Texte, Bilder und sonstige Inhalte in diesem Dokument unterliegen dem Schutz des Urheberrechts gemäß dem Urheberrechtsgesetz der Bundesrepublik Deutschland. Die Inhalte können von Ihnen nur dann genutzt und vervielfältigt werden, wenn Ihnen dies im Einzelfall durch den Rechteinhaber oder die Schrankenregelungen des Urheberrechts gestattet ist. Jede Art der Nutzung zu gewerblichen Zwecken ist untersagt. Zu den Möglichkeiten einer Lizenzierung von Nutzungsrechten wenden Sie sich bitte direkt an die verantwortlichen Herausgeberinnen/Herausgeber der entsprechenden Publikationsorgane oder an die Online-Redaktion des Deutschen Archäologischen Instituts (info@dainst.de). Etwaige davon abweichende Lizenzbedingungen sind im Abbildungsnachweis vermerkt.

Terms of use: By downloading you accept the terms of use (<https://publications.dainst.org/terms-of-use>) of iDAI.publications. Unless otherwise stated in the document, the following terms of use are applicable: All materials including texts, articles, images and other content contained in this document are subject to the German copyright. The contents are for personal use only and may only be reproduced or made accessible to third parties if you have gained permission from the copyright owner. Any form of commercial use is expressly prohibited. When seeking the granting of licenses of use or permission to reproduce any kind of material please contact the responsible editors of the publications or contact the Deutsches Archäologisches Institut (info@dainst.de). Any deviating terms of use are indicated in the credits.


Forschungen zur Archäologie
Außereuropäischer Kulturen
Band 8



MONGOLIAN - GERMAN KARAKORUM EXPEDITION

VOLUME 1

Excavations in the Craftsmen Quarter
at the Main Road



edited by
Jan Bemann, Ulambayar Erdenebat, & Ernst Pohl



F A A K 8

Kommission für Archäologie
Außereuropäischer Kulturen
des
Deutschen Archäologischen
Instituts, Bonn

DEUTSCHES
ARCHÄOLOGISCHES
INSTITUT



Kommission für
Archäologie
Außereuropäischer
Kulturen

in collaboration with

RHEINISCHE FRIEDRICH-
WILHELMS-UNIVERSITÄT
BONN

Vor- und Frühgeschichtliche Archäologie

MONGOLIAN
ACADEMY OF
SCIENCES

Institute of Archaeology


Forschungen
zur Archäologie
Außereuropäischer
Kulturen

Band 8

MONGOLIAN-
GERMAN
KARAKORUM
EXPEDITION VOLUME 1

Excavations in the
Craftsmen Quarter
at the Main Road

edited by
Jan Bemmann,
Ulambayar Erdenebat,
and Ernst Pohl



Reichert Verlag
Wiesbaden 2010

Bibliografische Informationen
der Deutschen Nationalbibliothek
Die Deutsche Nationalbibliothek
verzeichnet diese Publikation in der
Deutschen Nationalbibliografie; detaillierte
bibliografische Daten sind im Internet über
<http://dnb.d-nb.de> abrufbar

Copyright ©
Kommission für Archäologie
Außereuropäischer Kulturen 2010

(= Bonn Contributions to Asian Archaeology 2
Vor- und Frühgeschichtliche Archäologie der
Rheinischen Friedrich-Wilhelms-Universität Bonn)

Satz und Gestaltung:
Linden Soft Verlag e.K., Aichwald
Umschlaggestaltung: Hans-Peter Wittersheim
Druck und Vertrieb:
Reichert Verlag, Wiesbaden

ISBN 978-3-89500-697-5

All rights reserved.
No reproductions of any kind without
written permission.
Printed in Germany · Impreso en Alemania
Printed on fade resistant and
archival quality paper (PH 7 neutral) · tcf

Contents

Preface	7
1. W. SCHWANGHART, S. MÖLLER, B. SCHÜTT, Environmental Characteristics of the Upper and Middle Orkhon Valley, Mongolia	14
2. E. BECKER, Die sowjetisch-mongolischen Ausgrabungen von 1948/49	27
3. I. ELIKHINA, The Most Interesting Artefacts from Karakorum in the Collection of The State Hermitage Museum, St. Petersburg	39
4. U. ERDENEBAT, M. JANSSEN-KIM, E. POHL, Two Ceramic Deposits from the Territory of Karakorum	49
5. E. POHL, The Excavations in the Craftsmen-Quarter of Karakorum (KAR-2) between 2000 and 2005 – Stratigraphy and Architecture	63
6. E. POHL, KAR-2, Catalogue of Artefacts I: Processed Bones (with Faunal Analysis by A. von den Driesch and J. Peters)	137
7. E. POHL, KAR-2, Catalogue of Artefacts II: Spindle Whorls and other Clay Objects	165
8. G. NOMGUUNSÜREN, Preliminary Study of Cart Wheel-Bushings from Karakorum, Mongolia	173
9. CHR. LEE, Human Skeletal Remains from the Excavations in the Craftsmen-Quarter of Karakorum (KAR-2)	213
10. M. RÖSCH, E. FISCHER, T. MÄRKLE, B. OYUNTUYA, Medieval Plant Remains from Karakorum, Mongolia	219
11. A. VON DEN DRIESCH, J. PETERS, L. DELGERMAA, Animal Economy in the Ancient Mongolian Town of Karakorum. Preliminary Report on the Faunal Remains	251
12. K. KELM, Chemical Investigations on some Archaeological Findings from Karakorum	271
13. D. BAYAR, V. E. VOITOV, Excavation in the Islamic Cemetery of Karakorum	289
14. J. BEMMANN, E. POHL, B. SCHÜTT, W. SCHWANGHART, Archaeological Findings in the Upper and Middle Orkhon Valley and their Geographical Setup	307
15. U. B. BARKMANN, Die Geschichte des Klosters Erdeni Joo oder das Prinzip der Verflechtung von Staat und Religion	321
Supplements	

Wolfgang Schwanghart, Steffen Möller, Brigitta Schütt

Environmental Characteristics of the Upper and Middle Orkhon Valley, Mongolia

INTRODUCTION

The Orkhon Gol (Orkhon River) in central Mongolia is one of the most important drainages of the Khangai Mountains. Its upper and middle courses constitute not only a vein of present livelihoods but also represent major axes of migration and settlements during historic and prehistoric times (Bemmann et al., in this issue). Hence, various archaeological studies have been conducted in this area during the last decade.

Throughout time the environmental settings and their changes have always been a crucial factor in the rise and fall of human settlement dynamics. Thus, in order to understand the societal and cultural evolution in this area of interest detailed knowledge on its environmental controls and their spatio-temporal changes is required.

Here, we present an overview on the environmental characteristics of the Upper and Middle Orkhon Valley based on results gained during field campaigns between 2004 and 2006, environmental data analysis and literature reviews. Many environmental features, however, are only poorly documented in the “Western” literature mainly due to Mongolia’s remoteness and historical isolation from the rest of the world.

AREA OF INTEREST

The area of interest is located in the Arkhangai and Övörkhongai province. It encompasses the region between 47°00’ to 47°53’N and 102°15’ to 103°08’E and comprises elevations between 1210 m and 2350 m (Fig. 1–2). The term Upper Orkhon Valley refers to the part of the

river course located in the steeply intersected northern declivity of the Khangai Mountains where fluvial processes of bedrock detachment and incision prevail.

A sudden change to a predominance of gravel and sand deposition occurs where the Orkhon River exits the Khangai Mountains and forms the Middle Orkhon Valley (Fig. 2). Loosing its transport capacity due to rapid decrease in downward slope and, thus, flow velocity the river accumulated a vast fan. The medieval Mongolian capital Karakorum (and the modern Sum capital Kharkhorin) is located at the apex of this fan. From here the present river course meanders in the western part of the Middle Orkhon Valley until its confluence with Tamir River nearby Lake Ögii Nuur.

In the east of the Middle Orkhon Valley the Khögshin Orkhon River (Old Orkhon River) drains the eastern part of the Khangai Mountains and flows into Lake Ögii Nuur. Ögii Nuur is a quasi-endorheic freshwater lake – predominantly fed by the Khögshin River. It is only episodically an open basin during high water levels. During these phases the alluvial plain between the Orkhon River and the lake is flooded by lake water and drainage happens along Orkhon River (Fig. 2).

ENVIRONMENTAL CHARACTERISTICS

Mongolia’s exogenically driven environmental characters (e. g. climate, soils, vegetation and drainage) exhibit a strong zonal gradient determined by the planetary settings (Haase 1983). Thus, the mean annual precipitation sum decreases from 300–400 mm in the north to 100 mm and less in the south (Barthel 1983). Mainly governed by effective moisture supply




Fig. 1. Location of the area of interest (black outline). Elevation values are based on GTOPO30 (Projection: Eckert IV, Graticule: WGS84).

vegetation pursues this trend. Taiga forests in the north pass over into forest steppe followed by steppe, desert steppe and the Gobi desert in southward direction (Gunin et al. 1999).

Breaks in the zonal trend of the environmental settings may be caused by several local or regional peculiarities, however, the most important factor is elevation. In general, an increase in elevation causes a change in climate: with increasing elevation mean air temperature decreases while precipitation sums usually increase. The latitudinal extent of the study site is relatively low. However, parts of the area of interest are located in the Khangai Mountains. Hence, to characterize the environmental setting altitudinal zones have to be taken into account.

GEOLOGY

Geological maps of Mongolia reveal a variety and complexity of rock types and structures covering the time span from Precambrian to Quaternary. This diversity is due to successive terrane accretions to the Siberian Craton and consequent structuring during the Early and Late Palaeozoic. These structures were repeatedly reactivated during the Mesozoic

and Tertiary responding tectonic activity in western and central China, probably triggered by the Himalayan orogeny and Tibetan Plateau uplift (Traynor / Sladen 1995 and several authors therein).

The area of interest comprises several stratigraphical units from the Proterozoic and Cambrian (>50 Ma B.P.) to present. Oldest units can be found in the southeast and northwest of the Middle Orkhon Valley along a southeast-northwest stretching zone and composed of high grade metamorphic rock (mainly schists). Attached to this zone Devonian sediments are intermitted by plutonites and intrusions. The ridges of the Khangai Mountains are largely composed of Devonian to Mid-Permian sediments of marine facies that were subsequently orogenetically elevated, folded and ruptured in the response to compressional tectonics in southern Mongolia (Akademii Nauk 1990; Sladen / Traynor 2000; Walther 2005).

In the course of this tectonic activity but also during the Varisciden folding of the Khangai Mountains various plutonites and dikes formed and were exhumed by erosional processes during the Cenozoic. Induced by uplift widespread erosion and sedimentation processes in the mountain foreland and intramontane basins occurred from Triassic to




Fig. 2. First three principal components (PC) of Landsat ETM+ imagery (22 July 2002, R-PC1, G-PC2, B-PC3, Projection: UTM 48N, WGS84). Information in the second PC (green) can be correlated with vegetation.



Fig. 3. Landscape characteristics in the Upper and Middle Orkhon Valley. a) Exposition dependent *Larix siberica* in the northern declivity of the Khangai Mountains and the Middle Orkhon Valley; b) Basalts in the Upper Orkhon Valley; c) Tertiary weathering of granites in the Upper Orkhon Valley; d) Lake Ögii Nuur.

early Jurassic (Traynor / Sladen 1995). Rifting and synrift deposition resulted in the formation of an extensive belt of Late Jurassic to early Cretaceous basins especially in southern and eastern Mongolia (Sladen / Traynor 2000). Remnants of Jurassic thin layered, highly fissile lacustrine mudstones surrounding Ögii Nuur and in the Middle Orkhon Valley may be attributed to this period. In the vicinity of Ögii Nuur reddish conglomerates are exposed that can be attributed to the late Cretaceous and Tertiary and testify fluvial dynamics with high bedload transport as documented by pebble beds and channel bar sandstones (Traynor / Sladen 1995).

From Tertiary to present tectonic faulting was accompanied by intensive volcanism depositing flood basalts in the Upper Orkhon Valley and

in the east of Ögii Nuur (Walther 2005). Along the Middle Orkhon Valley thick deposits of Quaternary unconsolidated debris and fluvial pebble and sand accumulations predominate. Variability in discharge caused the forming of fluvial terraces (Fig. 4) traceable on both sides of the Middle Orkhon Valley. Glaciation only played a minor role during the Pleistocene and was limited to the elevations higher than 3000 m (Lehmkuhl / Lang 2001).

CLIMATE

Analysis and description of local or regional climatic characteristics is restricted by the scarce data availability regarding both temporal and spatial resolution. Moreover, data quality

is often poor. In order to characterize the regional climate data from the NOAA National Climate Data Center Global Historical Climate Network for the station Arvaiheer (1813 m) and Bulgan (1210 m) is used (Fig. 1).

According to the classification concept of Köppen (Kottek et al. 2006) the climate in the area of interest can be described as BSk climate in the south and as Dwc climate in the north. Climate is controlled by the extreme continentality and the large scale circulation patterns that produced a pronounced seasonal cycle of air temperature and precipitation. Annual precipitation totals 209 and 314 mm and mean annual temperature is ca. 1.5 and -1.4°C in Arvaiheer and Bulgan, respectively.

During the winter months the predominance of the Siberian High causes unhindered outgoing heat radiation and the subsequent cooling results in mean January air temperatures below -25°C. Minimum daily temperatures may even fall below -40°C where temperature inversions develop in basins and valleys (Barthel 1983). Due to the stable Siberian High precipitation during winter remains low and snow cover rarely exceeds 10 cm.

Prevailing west winds influence the Mongolian climate during the summer months. Monthly July air temperature averages around 15°C but maximum daily temperature may exceed 35°C. July is the wettest month of the year, too, but there is a strong interannual variation in monthly rainfall amounts (Fig. 5 b; 6 b) (Gunin et al. 1999). Rainfall is mainly short and intense and often linked to heavy thunderstorms that may occur several times per day. Data on rainfall duration is unavailable but own field observations suggest that rainfall events hardly exceed three hours.

Results of an estimation of recurrence intervals of daily precipitation amounts are shown in Fig. 5 c and 6 c. The daily rainfall magnitude-frequency relationship was modelled using the approach proposed by Ahnert (1986). Visually the logarithmic function fits the observational data well, however, for larger precipitation amounts (> 30 mm/day) there is an increase in uncertainty of the model due to scarce observational data for these extreme events. According to this model statistically every 1 (5, 10, 30) years a daily rainfall of 25 mm (38, 43, 52 mm) can be expected in Arvaiheer. It is assumed, however, that these values are underestimated because wind-induced undercatch, wetting loss and evaporation loss cause a precipitation amount bias that is es-




Fig. 4. Outcrop of the Orkhon terrace close to Kharkhorin exhibits fluvial deposits composed of basalt pebbles which are cryoturbatically reworked and covered by aeolian silts and sands.

pecially strong in forests and prairies (Zhang et al. 2004).

The transitional seasons spring and autumn are short and often characterized by strong temperature increases and decreases. During this time frequent occurrence of freezing-thawing cycles is an important factor for physical weathering processes (Fig. 5 a; 6 a). Due to the ambiguous correlation of the diurnal temperature range and elevation (Geerts 2003) extrapolation of these results to other altitudinal zones is not possible, yet.

The harsh climate conditions entails in many constraints for agricultural activity. Singular weather events, however, even aggravate the problems especially for livestock owners. These events are termed 'Zuds' and belong to the most dramatic catastrophes leading to high economic losses for herders. In general Zuds are caused by a sequence of extreme weather conditions




Fig. 5. Analysis of daily climate station data of Avaiheer during 1960/1961 to 1998. (a) Seasonal development of mean maximum and minimum daily temperatures and average number of days with maximum temperatures greater than 0°C and minimum temperatures less than 0°C (freezing-thawing cycles). (b) Monthly precipitation amounts and variability. (c) Recurrence intervals of daily precipitation amounts.




Fig. 6. Analysis of daily climate station data of Bulgan during 1960/1961 to 1998. (a) Seasonal development of mean maximum and minimum daily temperatures and average number of days with maximum temperatures greater than 0°C and minimum temperatures less than 0°C (freezing-thawing cycles). (b) Monthly precipitation amounts and variability. (c) Recurrence intervals of daily precipitation amounts.



Fig. 7. Kastanozem profile developed in a top to bottom sequence of colluvial/alluvial deposits, aeolian fines and slope debris. Below 60 to 70 cm a indurated calcrete horizon has established.

which in combination cause unavailability of nourishments or water for the herds. Starvation and dehydration of the animals during a devilish Zud winter in 1944/45 caused a nationwide livestock reduction of approx. 30%. Two consecutive Zud disasters in 1999–2000 and 2000–2001 resulted in a livestock loss of 29,640 animals in the Arkhangai province only (Goroncy 2006).

SOILS

As in most parts of Mongolia loamy and sandy soils prevail in the area of interest, usually riddled with a considerable amount of detritus or pebbles (Gunin et al. 1999). Besides, hydromorphic soils occur in the vicinity of rivers and lakes. In the slightly rolling hills of the Middle Orkhon Valley and in intra-montane basins there is a predominance of Kastanozems in regions below 1700–2000 m. (Haase 1983). On steep slopes these steppe soils are poorly developed, frequently only

a few decimetres thick and formed in debris cover with sandy to loamy matrix (weathered layer). Much thicker weathered layers can be found in the foothills and valley floors. Parent material of these soils is rich in skeleton which is often layered and shows adjusted pebbles and debris (Fig. 7). It is assumed that an interfingering of autochthonous weathered material and soil sediments (colluvial deposits) prevails in the locations. Usually calcrete layers and chalk precipitated on the bottom of debris and pebbles are found in variable depths in the Kastanozem zone (Fig. 7) (Haase 1983; Klinge 2001) wherever the hydrologic gradient is small enough to allow ascendent soil water movement. Caragana roots penetrating the soil up to two meters demonstrate strong soil rooting.

In the mountainous areas a differentiation of the Kastanozems occurs according to slope exposition. Due to a stronger radiation and subsequent increased evaporation soils tend to be drier and lighter in south- and westward inclined slopes while north- and eastward inclined slopes have more favourable moisture conditions (Haase 1983; Klinge 2001).

Between 1700 and 2400 m more humid conditions and stronger exposition to solar radiation lead to a development of A–C-soils with thick humus horizons due to a lack of decomposition of litter (Paratschernosems). Permafrost and the extremely low winter temperatures prevent the dislocation of soluble and promote podzol formation producing patches of Cryotaigasols (Haase 1983).

Above 2000 to 2400 m partly thick (up to 80 cm) meadow soils prevail. They are characterized by a distinct humus-rich top soil (Dernosem) above a C-horizon (A–C-soil). Their strong relationship to steppe soils constitutes in high humus contents up to 8 mass-%. They interfinger with peat bogs that develop close to springs and in topographic sinks where ground ice impedes percolation (Haase 1983).

Above 2900 to 3200 m debris and boulders prevail and a development of soil is constrained due to cryoturbatic processes and the lack of vegetation.

VEGETATION

The general vegetation pattern of Mongolia follows a zonal, climate dependent sequence (Gunin et al. 1999). Transitions between these

zones are rather continuous but often constitute in exposition-depending vegetation mosaics (Hilbig 2000; Dulamsuren et al. 2005b).

The major part of the study site is situated in the steppe region. The higher elevated regions of the Khangai Mountains can be classified as mountain steppe or mountain forest steppe (Barthel et al. 1962). Here, sparse *Larix*-populations are mainly restricted to northern exposed slopes above 1350–1400 m (Fig. 3 a) (Walther / Gegeensuvd 2005). The sharp contrast to the deforested southern exposed slopes in the mountainous regions is attributed to the different exposition to solar radiation causing favourable moisture, temperature and edaphically conditions (Treter 1996; Sommer / Treter 1999). Along the Orkhon River locally riverine forests assembled predominantly of *Ulmus* and *Salix* occur. The slightly rolling plains of the Middle Orkhon Valley are covered by steppe type vegetation which can be subdivided into the three major types *Artemisia*-, *Stipa*- and *Allium*-Steppe.

The reasons for the present dominance of steppe vegetation in the area of interest are disputed. Extremely high grazing pressure and unregulated lumbering suggest that the present day vegetation is strongly influenced by human activity since earlier than the Medieval (Hilbig 2000; Rösch et al. 2006). Fernandez-Gimenez / Allen-Diaz (2001) and Möller et al. (2005) show that livestock influences plant species composition by defoliation, trampling, nutrient enrichment and depletion over the broader landscape scale. Dulamsuren et al. (2005a), however, doubt the general significance of human activity for producing the north-south pattern of vegetation units in the forest-steppe ecotone by presenting evidence that site conditions like moisture supply and radiation vary in a short spatial range and determine ground vegetation. Furthermore, it has been shown that repeated expansion and retreat of steppe vegetation correlated with decreasing and increasing precipitation since 15 ka B.P. (Tarasov et al. 2000). Presumably both human activity and climate govern present vegetation dynamics (Gunin et al. 1999).

DRAINAGE

The major drainages are perennial (Orkhon River, Khögshin and Tamir River). Due to highly mobile channels these rivers can be classified as braided in the vast Middle Orkhon




Fig. 8. Channel morphology at Ögii Nuur headwater area. The picture was taken one day after a heavy rainfall event. The site shown has a drainage area of around 0.5 km².

Valley. Following the climatic seasonality in precipitation the Orkhon discharge has its maximum during July with a monthly average around 100 m³/s (<http://www.sage.wisc.edu/riverdata>, 10. Dec. 2006).

Episodic run-off during high precipitation events results in the incision of channels into the thick debris cover and locally into bedrock (Fig. 8). These channels, however, are rarely connected and do not form a regional drainage network. They are arheic where they enter broader valleys or flat to slightly inclined terrain. In the catchment adjacent to Ögii Nuur it was observed that concentrated surface run-off during high rainfall events rarely reaches the lake but prior infiltrates. Hence, it is suggested that these drainage characteristics are a result of a factor combination of high infiltration in sandy and silty soils and short but intensive rainfall events.

GEOMORPHOLOGICAL AND PALAEOENVIRONMENTAL IMPLICATIONS

The environmental characteristics of the Upper and Middle Orkhon Valley presented are the results of endogenic and exogenic processes that vary in space and time. These processes create a huge diversity of materials and landforms that differ in scale and life time (Ahnert 1996). Thus, landforms and their associated materials offer archives to be analyzed in order to understand the underlying processes of their genesis. Since many of these landforms are relic or inactive they allow insight into palaeoenvironments and their dating enables their temporal placement. Selected examples of landforms shall clarify this issue in terms of the area of interest.

LAKE SYSTEMS

The water input-output ratio of lakes is an important measure to evaluate the climatic evolution of catchments. Closed-basin lakes react on changes of this ratio by increasing or decreasing water levels and establish a shoreline when the water level restabilizes at a new position (Sack 2001; Street-Perrott 1980). Depending on the substratum and speed and type of lake level change altering shorelines can be deduced from beach ridges, beach terraces or cliff edges (Street-Perrott / Harrison 1985). In Mongolia and north-western China these landforms have proven to be valuable indicators for palaeoenvironmental interpretation (e.g. Grunert et al. 2000; Pachur et al. 1995; Walther 1998; 1999; Walther et al. 2003; Wünnemann et al. 1998; Wünnemann / Hartmann 2002). In the Upper and Middle Orkhon Valley only a few lakes exist with Ögii Nuur being the largest. Walther / Gegeensuvd (2005) describe for Ögii Nuur the existence of beach ridges in 2.5 and 3.5 m above the present lake level. Yet, as Ögii Nuur is only episodically a closed lake basin and opens in the west towards the Orkhon River, the question on the damming of the lake remains unexplained. Smaller lakes in the Orkhon catchment, however, show strong interannual fluctuations in lake level. Landsat imagery available for 1995 and 1999 show closed water surfaces in the Middle Orkhon Valley close to Ögii Nuur Sum that have vanished in 2001 and 2002.

Besides geomorphological evidences lake systems offer lacustrine sediments that are

the definite deposition in a complex storage system and, thus, constitute an answer to the drainage basin's geomorphological reaction to climatic, tectonic or human impacts. The continuity of lacustrine sediments is their big advantage if discrete spatial information is lacking. The signal given by lacustrine sediments is mixed, allowing conclusions on the paleolimnic environment (diatoms, authigenic minerals), paleotemperature (oxygen isotopes of shells and authigenic minerals), the local to regional vegetation cover (pollen) and the overall weathering conditions in the drainage basin (bulk chemistry, clay minerals).

In addition to the deposition of allochthonous detritus, lacustrine sediments are characterized by precipitation of authigenic minerals from aqueous solutions. Precipitation of authigenic minerals is substantially influenced by lake water salinity and its chemical composition, thus, authigenic minerals are particularly of interest for the reconstruction of the palaeolimnic environments. In contrast, the pedogenic minerals (such as clay minerals, oxides) in lacustrine sediments contain information about the synsedimentary environmental conditions in the drainage basin (Schütt 1998; 2004).

Lacustrine sediments extracted from Ögii Nuur have been analyzed on organic matter (total organic carbon [TOC]), diatoms and pollen distribution and provide evidence for varying lake levels and moisture supply during the Holocene (Walther / Gegeensuvd 2005). Pollen profiles of core UGI-3 were analyzed by Röscher et al. (2006). They suggest increased woodland distribution during the middle Holocene caused by increased precipitation. A subsequent decline of the arbor-nonarbor pollen ratio is interpreted as human induced (Röscher et al. 2006). Further analysis on mineralogical and chemical parameters is currently implemented.

RELIEF AND MORPHODYNAMICS

Information on drainage basin geomorphodynamics, on the predominating processes of detrital material flows, magnitude and frequency of these processes and their climatic, tectonic or anthropogenic impulses can only be poorly deduced from lacustrine sediments. In contrast, this information is given by temporally discontinuous, ephemeral sediments to be found in the drainage basin, such as flu-

vial and colluvial deposits (e.g., alluvial fans, river terraces), sediments of mass movements etc. Although these sedimentary archives lack temporal continuity, they have the advantage of being spatially discrete.

Periglacial landforms and their spatial occurrence are strongly associated with climate. As afore mentioned there is a strong climatic dependence on elevation that is reflected by the spatial distribution of soils and vegetation. Pękala / Repelewska-Pękalowa (1993) and Lehmkuhl / Lang (2001) determine the present lower limit of active cryoplanation terraces and solifluction forms at 2700–2900 m in the Khangai Mountains. Discontinuous permafrost and smaller but active forms are found as low as 2200 m on northern exposed slopes. Findings of inactive ice-wedges as low as 1400 m, however, suggest significant changes of temperature and precipitation during the Pleistocene in the Khangai Mountains (Lehmkuhl / Lang 2001) that have been previously recognized by Owen et al. (1997) and Owen et al. (1998) in the Gobi Desert. Glaciation during the Last Glacial Maximum is likely to have never occurred in the area of interest. A depression of the snow line or equilibrium-line altitude (ELA) to 2700–2800 m in the Khangai Mountains is suggested by Lehmkuhl et al. (2004) and Lehmkuhl / Lang (2001) which did not affect the area of interest.

Aeolian sediments play an important role in landscape evolution especially during arid periods. In Mongolia they are widespread and occur independently from altitude. Their deposition happened both during the Pleistocene glacial periods and interstadials (Grunert / Klein 1998; Grunert / Dasch 2004; Grunert / Lehmkuhl 2004). However, meteorological observations of dust storms provide evidence for present aeolian activity (Natsagdorj et al. 2003; Dulam 2005). Reworked fluvial sediments of large rivers and dried out lakes are identified as sources of the wind blown material (Grunert / Lehmkuhl 2004). In the Upper and Middle Orkhon Valley several dune fields or sand sheets exist, often vegetated by *Gramineae achnatherum*. Surfaces with strong aeolian accumulation have been identified by a supervised classification of Landsat ETM+ imagery (cf. Bemmman et al. in this issue, Fig. 2). The identified regions are located in the vicinity of rivers and in the footzones of hills. Due to the prevalence of WNW wind directions a preponderant occurrence of aeolian sand sheets on south and southeast facing slopes may indicate an

orographically induced settling in the wind shadow of mountains.

PERSPECTIVE

Cryoturbatic disturbance of periglacial debris, accumulation of material displaced by sheet flow, interfingering fans and aeolian loess-like sedimentation create a relatively uniform geomorphological appearance of terrestrial sedimentary architectures despite their variable processes of origin (Haase 1983). This uniformity makes it a challenge to unravel the complexity of landscape evolution. An ongoing human interference with landscape forming processes since prehistoric times (Rösch et al. 2006) even complicates this process system. Changes in human activities like lumbering, animal husbandry and cultivation of grains affect the environmental settings and modify geomorphological process intensities. In order to address the issue of landscape evolution and to understand the interactions with cultural and societal development in the Upper and Middle Orkhon Valley an integrated approach combining physio-geographical and archeological research is required and promising.

Addresses:

Dipl. Geogr. Wolfgang Schwanghart
Universität Basel, Departement Umweltwissen-
schaften, Physiogeographie und Umweltwandel
Klingelbergstrasse 27
CH-4056 Basel

w.schwanghart@unibas.ch

Dr. Steffen Möller
Georg-August-Universität Göttingen,
Geographisches Institut
Goldschmidtstraße 5
D-37077 Göttingen

steffen.moeller@geog.uni-goettingen.de

Prof. Dr. Brigitta Schütt
Freie Universität Berlin, Institut für Geographische
Wissenschaften, Physische Geographie
Maltheserstr. 74–100, Haus H
D-12249 Berlin

schuett@geog.fu-berlin.de

Figure credits:

1–3, 5–6, 8 Schwanghart; 4, 7 Möller.

BIBLIOGRAPHY

- Ahnert, Frank
1986 The magnitude-frequency index – an approach to the identification of hygric morphoclimates. In: Zeitschrift für Geomorphologie, Suppl.-Band 60: 1–6. Berlin.
- 1996 Einführung in die Geomorphologie. Stuttgart.
- Akademiia Nauk
1990 Mongol'skaia Narodnaia Respublika Natsional'nii Atlas. Ulaanbataar.
- Barthel, Hellmuth
1983 Die regionale und jahreszeitliche Differenzierung des Klimas in der Mongolischen Volksrepublik. In: Barthel, Hellmuth / Brunner, Horst / Haase, Günter (eds.), Physisch-Geographische Studien in Asien. *Studia Geographica* 34: pp. 3–91. Brno.
- Barthel, Hellmuth / Haase, Günter / Richter, Hans
1962 Die mongolische Volksrepublik. In: Zeitschrift für den Erdkundeunterricht 14: 288–318. Berlin.
- Dulam, Jugder
2005 Discriminate analysis for dust storm prediction in the Gobi and steppe regions in Mongolia. In: Water, Air and Soil Pollution 5: 37–49. Dordrecht.
- Dulamsuren, Choimaa / Hauck, Markus / Muehlenberg, Michael
2005a Ground vegetation in the Mongolian taiga forest-steppe ecozone does not offer evidence for the human origin of grasslands. In: *Applied Vegetation Science* 8: 149–154.
- 2005b Vegetation at the taiga forest-steppe borderline in the western Khentey Mountains, northern Mongolia. In: *Annales Botanici Fennici* 42: 411–426. Helsinki.
- Fernandez-Gimenez, Maria / Allen-Diaz, Barbara
2001 Vegetation change along gradients from water sources in three grazed Mongolian ecosystems. *Plant Ecology* 157: 101–118. Dordrecht.
- Geerts, Bert
2003 Empirical estimation of the monthly-mean daily temperature range. In: *Theoretical and Applied Climatology* 74: 145–165. Wien.
- Goroncy, Florian
2006 Räumliche und zeitliche Analyse von Klimaelementen in Zentralasien für die Erklärung der Genese von Zud-Ereignissen in der Mongolei. Diploma-Thesis, Freie Universität Berlin.
- Grunert, Jörg / Dasch, D.
2004 Dynamics and evolution of dune fields on the northern rim of the Gobi Desert (Mongolia). In: Zeitschrift für Geomorphologie, Suppl.-Band 133: 81–106. Berlin.
- Grunert, Jörg / Klein, Michael
1998 Binnendünen im nördlichen Zentralasien (Uvs Nuur, westliche Mongolei). In: Böse, Margot / Hofmann, Jürgen (eds.), Forschungsergebnisse zur Klimageschichte und Reliefentwicklung Nordafrikas und Asiens, *Berliner Geographische Abhandlungen* 63: pp. 45–66. Berlin.
- Grunert, Jörg / Klein, Michael / Stumböck, Max / Dasch, D.
1999 Bodenentwicklung auf Altdünen im Uvs Nuur Becken. In: *Die Erde* 130: 97–116. Berlin.
- Grunert, Jörg / Lehmkuhl, Frank
2004 Aeolian sedimentation in arid and semi-arid environments of Western Mongolia. In: Smykatz-Kloss, Werner / Felix-Henningsen, Peter (eds.), *Paleoecology of Quaternary Drylands*. In: *Lecture Notes in Earth Sciences* 102: 195–218. Berlin, New York, Toronto.
- Grunert, Jörg / Lehmkuhl, Frank / Walther, Michael
2000 Paleoclimatic evolution of the Uvs Nuur basin and adjacent areas (Western Mongolia). In: *Quaternary International* 65–66: 171–192. Oxford.
- Gunin, Peter D. / Vostokova, Elizabeth A. / Dorofeyuk, Nadezhda I. / Tarasov, Pavel E. / Black, Clanton C.
1999 *Vegetation Dynamics of Mongolia*. *Geobotany* 26. Dordrecht, Boston, London.
- Haase, Günter
1983 Beiträge zur Bodengeographie der Mongolischen Volksrepublik. In: Barthel, Hellmuth / Brunner, Horst / Haase, Günter (eds.), *Physisch-Geographische Studien in Asien*. In: *Studia Geographica* 34: 231–367. Brno.

- Hilbig, Werner
2000 Forest distribution and retreat in the forest steppe ecotone of Mongolia. In: Marburger Geographische Schriften 135: 171–187. Marburg.
- Klinge, Michael
2001 Glazialgeomorphologische Untersuchungen im Mongolischen Altai als Beitrag zur jungquartären Landschafts- und Klimageschichte der Westmongolei. Aachener Geographische Arbeiten 35. Aachen.
- Kottek, Markus / Grieser, Jürgen / Beck, Christoph / Rudolf, Bruno / Rubel, Franz
2006 World Map of the Köppen-Geiger climate classification updated. In: Meteorologische Zeitschrift 15: 259–263. Stuttgart.
- Lehmkuhl, Frank / Klinge, Michael / Stauch, Georg
2004 The extent of Late Pleistocene glaciations in the Altai and Khangai Mountains. In: Ehlers, Jürgen / Gibbard, Philip. L. (eds.), Quaternary glaciations extent and chronology. Part III: South America, Asia, Africa, Australasia, Antarctica. Developments in Quaternary Science Vol. 2: pp. 243–254. Amsterdam, Boston, Heidelberg.
- Lehmkuhl, Frank / Lang, Andreas
2001 Geomorphological investigations and luminescence dating in the southern part of the Khangai and the Valley of the Gobi Lakes (Central Mongolia). In: Journal of Quaternary Science 16 (1): 69–87. Chichester.
- Möller, Steffen / Schütt, Brigitta / Walther, Michael
2005 Land degradation and soil erosion in the Kharkhorin region, Republic of Mongolia. In: Proceedings of the Third International Congress on Environmental Change in Central Asia, pp. 86–91. Ulaanbaatar.
- Natsagdorj, Luvsan / Dulam, Jugder / Chung, Yong-Seung
2003 Analysis of dust storms observed in Mongolia during 1937–1999. In: Atmospheric Environment 37: 1401–1411. Oxford.
- Owen, Lewis A. / Richards, Ben / Rhodes, Edward J. / Cunningham, W. Dickson / Windley, Brian F. / Badamgarav, J. / Dorjnamjaa, D.
1998 Relic permafrost structures in the Gobi of Mongolia: age and significance. In: Journal of Quaternary Science 13: 539–547. Chichester.
- Owen, Lewis A. / Windley, Brian F. / Cunningham, W. Dickson / Badamgarav, J. / Dorjnamjaa, D.
1997 Quaternary alluvial fans in the Gobi of southern Mongolia: evidence for neotectonics and climate change. In: Journal of Quaternary Science 12: 239–252. Chichester.
- Pachur, Hans J. / Wünnemann, Bernd / Zhang, Hucai
1995 Lake Evolution in the Tengger Desert, Northwestern China, during the Last 40000 Years. In: Quaternary Research 44 (2): 171–180. New York.
- Pękala, Kazimierz / Repelewska-Pękalowa, Janina.
1993 Solifluction and related slope processes in the mountains of Central Asia (Mongolia). In: Frenzel, Burkhard (ed.): Palaeoclimate Research, Special Issue: ESF Project “European Palaeoclimate and Man” 6/11: pp. 87–101. Stuttgart.
- Rösch, Manfred / Fischer, Elske / Märkle, Tanja
2006 Human diet and land use in the time of the Khans – Archaeobotanical research in the capital of the Mongolian Empire, Karakorum, Mongolia. In: Vegetation History and Archaeobotany 14: 485–492. Berlin/Heidelberg.
- Sack, Dorothy
2001 Shoreline and basin configuration techniques in paleolimnology. In: Last, William M. / Smol, John P. (eds.), Tracking environmental change using lake sediments. Vol. 1: Basin analysis, coring and chronological techniques, pp. 49–71. Dordrecht.
- Schütt, Brigitta
1998 Chemical and mineralogical characters of lacustrine sediments as paleoenvironmental indicators – An example from the Laguna Jabonera, Central Ebro Basin. In: Terra Nostra 98/6: 115–120. Berlin.
- 2004 Reconstruction of Holocene weathering conditions from chemical character of playa-lake-sediments – a case study from central Spain. In: Werner Smykatz-Kloss / Peter Felix-Henningsen (eds.), Paleoeecology of Quaternary Drylands. Vol. 102 of Lecture Notes in Earth Sciences: pp. 5–30. Berlin, New York, Toronto.

- Sladen, Chris / Traynor, James J.
2000 Lakes during the evolution of Mongolia. In: Gierlowski-Kordesch, Elizabeth H. / Kelts, Kerry R. (eds.), *Lake basins through space and time. American Association of Petroleum Geologists Studies in Geology* 46: pp. 35–57. Tulsa.
- Sommer, Martin / Treter, Uwe
1999 Die Lärchenwälder der Gebirgswaldsteppe in den Randgebirgen des Uvs Nuur-Beckens. In: *Die Erde* 130: 173–188. Berlin.
- Street-Perrott, F. Alayne
1980 The relative importance of climate and hydrogeological factors in influencing lake level fluctuations. In: *Palaeoecology of Africa* 12: 137–158. Rotterdam.
- Street-Perrott, F. Alayne / Harrison, Sandy P.
1985 Lake levels and climate reconstructions. In: Hecht, Alfred D. (ed.), *Palaeoclimate analysis and modelling*, pp. 291–340. New York.
- Tarasov, Pavel E. / Dorofeyuk, Nadezhda / Metel’ Tseva, Elizaveta
2000 Holocene vegetation and climate changes in Hoton-Nur basin, north-west Mongolia. In: *Boreas* 29: 117–126. Oxford.
- Traynor, James J. / Sladen, Chris
1995 Tectonic and stratigraphic evolution of the Mongolian People’s Republic and its influence on hydrocarbon geology and potential. In: *Marine and Petroleum Geology* 12: 35–52. Amsterdam.
- Treter, Uwe
1996 Gebirgs-Waldsteppe in der Mongolei. In: *Geographische Rundschau* 48 (11): 655–661. Braunschweig.
- Walther, Michael
1998 Paläoklimatische Untersuchungen zur jungpleistozänen Landschaftsentwicklung im Changai-Bergland und in der nördlichen Gobi (Mongolei). In: *Petermanns Geographische Mitteilungen* 142 (3/4): 207–217. Gotha.
- 1999 Befunde zur jungquartären Klimaentwicklung rekonstruiert am Beispiel der Seespiegelstände des Uvs Nuur-Beckens (NW-Mongolei). In: *Die Erde* 130: 131–150. Berlin.
- 2005 Ein idealer Ort für ein festes Lager. Zur Geographie des Orchontals und der Umgebung von Charchorin (Karakorum). In: *Dschingis Khan und seine Erben. Das Weltreich der Mongolen*, pp. 128–132. München.
- Walther, Michael / Gegeensuvd, S.
2005 Ugii Nuur (Zentral Mongolei). Paläoumweltstudien zu Seespiegelschwankungen und zur holozänen Klimaentwicklung. *Geographica-Oekologica. Journal of MO-LARE Research Centre*, 2. Ulaanbaatar.
- Walther, Michael / Wünnemann, Bernd / Tschimek-saichan, Altangerel
2003 Seen und Paläoseen in der Mongolei und Nordwestchina. In: *Petermanns Geographische Mitteilungen* 147 (5): 40–47. Gotha.
- Wünnemann, Bernd / Hartmann, Kai.
2002 Morphodynamics and Paleohydrography of the Gaxun Nur Basin, Inner Mongolia, China. *Zeitschrift für Geomorphologie, Suppl.-Band* 126: 147–168. Berlin.
- Wünnemann, Bernd / Pachur, Hans J. / Li, J.
1998 Chronologie der pleistozänen und holozänen Seespiegelschwankungen des Gaxun Nur / Sogo Nur und Baijian Hu, Innere Mongolei, Nordwestchina. In: *Petermanns Geographische Mitteilungen* 142 (3/4): 191–206. Gotha.
- Zhang, Yinsheng / Ohata, T. / Yang, Daqing / Davaa, G.
2004 Bias correction of daily precipitation measurements for Mongolia. In: *Hydrological Processes* 18: 2991–3005. Chichester.